


FIG. 1 IC₅₀ values

| | Aromatic bpVs | | Polar bpVs | |
|-----------|---------------|----------|------------|----------|
| inhibitor | bpV | bpV | bpV | bpV |
| | (bipy) | (phen) | (HOpic) | (pic) |
| РТР-β | 60.3 nM | 343 nM | 4.9 μM | 12.7 μM |
| | +/- 9.6 | +/- 88.5 | +/- 0.9 | +/- 3.2 |
| PTP-1B | 164 nM | 920 nM | 25.3 μM | 61 μM |
| | +/- 22.6 | +/- 45.2 | +/- 2.9 | +/- 10.5 |
| PTEN | 18 nM | 38 nM | 14 nM | 31 nM |
| | +/- 0.8 | +/- 2.4 | +/- 2.3 | +/- 1.7 |











SUBSTITUTE SHEET (RULE 26)


5 / 10











10 / 10

low nanomolar inhibitors high µmolar aromatic nanomolar μ molar μ molar μ molar μ molar phosphorylation high nanomolar phosphorylation low nanomolar polar inhibitors high *µ*molar nanomolar stimulates μ molar μ molar μ molar inhibits in vivo actin in vitro pTyr in vitro Ly294002 (pTyr) inhibitor class pSer473(PKB) rapamycin (PKB) Inhibitor sensitivity PTEN inhibition Insulin mimetic PTP inhibition Cytotoxicity property

SUBSTITUTE SHEET (RULE 26)

FIG. 7